

October 2017 | Fact Sheet

## The HPV Vaccine: Access and Use in the U.S.

The human papilloma virus (HPV) vaccine is the first and only vaccination that helps protect women and men from getting many different types of cancer that are associated with different HPV strains. The vaccine protects young people against infection from certain strains of the HPV, the most common sexually transmitted infection (STI) in the United States.<sup>1</sup> Since HPV vaccines were first introduced in the U.S. in 2006 there have been changes in the range of protection they offer and the dosing regimen. Furthermore, the vaccines were originally recommended only for girls and young women and were subsequently broadened to include boys and young men. This factsheet discusses HPV and related cancers, use of the HPV vaccines for both females and males, and insurance coverage and access to the vaccines.

### HPV AND CANCER

There are more than 150 strains of HPV, and while most cases of HPV infection usually resolve on their own, there are more than 40 strains that can cause cancer. HPV infection in the U.S. is widespread; about 79 million Americans are currently infected with HPV and there are more than 14 million new infections annually.<sup>2</sup> More than 43% of American adults ages 18-59 are infected with genital HPV, with higher rates among men (45%) than women (40%).<sup>3</sup> Almost 50% of new infections occur in adults ages 15-24. HPV is related to almost 100% of cervical cancer cases, with two strains (16 and 18) related to approximately 70% of cervical cancer cases.<sup>4</sup> While cervical cancer is the main concern with HPV, the infection affects both women and men and is also known to cause oral, anal, vulvar, vaginal and penile cancers, as well as genital warts.<sup>5</sup>

### CERVICAL CANCER


Most cases of cervical cancer occur in women in developing countries, but it remains a challenge in the U.S. In the U.S., it is estimated that over 12,820 new cervical cancer cases will be diagnosed in 2017.<sup>6</sup> While cervical cancer is usually treatable, especially when detected early through routine screening with Pap tests, more than 4,210 deaths from cervical cancer will occur in 2017. Guidelines by the U.S. Preventive Services Task Force (USPSTF) recommend that most women ages 21 to 65 receive a Pap test once every three years.<sup>7</sup> In recent years, the USPSTF and some professional associations and committees have added a recommendation for screening certain populations of women with a test for high-risk HPV DNA.

#### U.S. Preventive Services Task Force Cervical Cancer Screening Recommendation

The [USPSTF](#) recommends screening for cervical cancer in women age 21 to 65 years with cytology (Pap smear) every 3 years or, for women age 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years.

- Despite widespread availability of pap testing, disparities in cervical cancer incidence and mortality rates by race persist. Black women have the highest mortality rates of the disease (**Figure 1**). More than half of cervical cancer cases are detected in women who have never been screened or have not been screened as frequently as recommended in guidelines.<sup>8</sup>

- One notable paradox however, is that Black women also have the highest rates of recent pap testing to screen for the disease (75%, compared to 69% of White women), yet experience the



highest mortality rate.<sup>9</sup> Limited access to treatment and diagnosis at later stages of disease progression, as well as cost, lack of physician referral, and cultural barriers may account for some of the disproportionate impact of cervical cancer on Black women.<sup>10</sup>

## THROAT CANCER

- Approximately 16,479 cases of throat cancer (oropharyngeal) occur annually in the U.S, most (70%) of which are associated with oral HPV infection, making it the second most common HPV-related cancer.<sup>11</sup> However, it's important to note that smoking and alcohol use are also risk factors. Oropharyngeal cancer is more common among men than women.<sup>12</sup> Research suggests that HPV vaccines can help protect against throat cancer, since many are associated with HPV 16, one of the strains that the vaccines protect against.<sup>13</sup>

## ANAL CANCER

- HPV is also responsible for the vast majority (95%) of the 8,200 annual cases of anal cancer in the U.S.<sup>14</sup> Most cases of anal cancer are among women, but men who have sex with men are also at higher risk.<sup>15</sup> HPV strains 16 and 18, as well as a history of cervical cancer, and suppressed immune systems are all risk factors for anal cancer.

## HPV VACCINES

Beginning in 2017, Gardasil<sup>®</sup>9 is the only HPV vaccine available in the U.S.


- The FDA first approved first-generation Gardasil<sup>®</sup>, produced by Merck, in 2006, which prevented infection of four strains of HPV – 6, 11, 16, and 18. In December 2014, Gardasil<sup>®</sup>9 was approved by the FDA. This vaccine protects against 9 strains of HPV: the four strains approved in the previous Gardasil vaccine, as well as 31, 33, 45, 52, and 58.<sup>16</sup>
- These strains are associated with the majority of cervical cancer, anal cancer, and throat cancer cases as well as most genital warts cases and some other HPV-associated ano-genital diseases.<sup>17, 18</sup>

- Gardasil®9 has been approved by the FDA for use in males and females ages 9-26. The federal Advisory Committee on Immunization Practices (ACIP) recommends that all girls and boys get vaccinated at age 11 or 12, and that girls and women ages 13-26 and boys and men ages 13-21 be given a “catch-up” vaccination.<sup>19</sup> The vaccine is recommended for use in men ages 22-26 if they have not been previously vaccinated, are immunocompromised, or engage in sexual activity with other men. ACIP recommended the vaccine for females in 2006 and added the recommendation for males in 2011.
- ACIP now recommends that 11-14 year olds receive two doses of the HPV vaccine over 6 months, instead of the original recommendation of three doses. Teens and young adults who initiate vaccination at age 15 through 26 should continue to receive three doses over six months.<sup>20</sup>
- These recommendations are designed to promote immunization when the vaccine is most effective – before the initiation of sexual activity and exposure to HPV. Those already infected with HPV also can benefit from the vaccine because it can prevent infection against HPV strains they may not have contracted, but the vaccine does not treat existing HPV infections. Additionally, the vaccine elicits a higher immune response from adolescents ages 11 to 12 than in older teens.<sup>21</sup>
- While the recommendation does not include women above age 26, some evidence supports providing HPV vaccines to adult women on a case-by-case basis.<sup>22</sup> Cost effectiveness of providing the vaccine to women over age 26 tends to decrease because women over the age of 26 are more likely to have been exposed to the specific serotypes and have fewer sexual partners and hence less risk of exposure. However, other countries, such as Australia, recommend the HPV vaccine for women through age 45.<sup>23</sup>
- Current research suggests the vaccine protection is long-lasting: more than 10 years of follow-up data indicate the vaccines are still effective and there is no evidence of waning protection, although it is still unknown if recipients will need a booster.<sup>24</sup>

## OUTREACH AND UTILIZATION

Awareness of the importance of the HPV vaccine has grown, and take-up has been increasing since it entered the market.

- In 2016, half of adolescent girls aged 13-17 were up-to-date with their HPV vaccinations (HPV UTD) (**Figure 2**).<sup>25</sup> Adolescent girls whose families live below the poverty line and girls living in urban areas were more likely to be HPV UTD compared to adolescent girls whose families live above the poverty line or live in rural areas. HPV vaccination rates among teen boys are lower than for girls (38% HPV UTD in 2016), but they have been rapidly rising since 2011.<sup>26</sup>



- Some people begin the vaccine series but do not complete it. In 2016, 65% of adolescent girls and 56% of boys received at least one dose of the HPV vaccine.<sup>27</sup>

- HPV vaccination rates vary by state, from 31% of adolescent girls being HPV UTD in South Carolina to 73% in Rhode Island (**Figure 3**).

- Presently at least 25 states and D.C. have laws that either require HPV vaccination for school entry, provide funding to cover the costs of the vaccines, or support public education about HPV and the vaccine. D.C. and Virginia require the vaccine for girls to enter the sixth grade, but allow parents to opt out of the requirement due to medical, moral, or religious opposition.<sup>28</sup> Rhode Island requires all seventh-grade students to be vaccinated.<sup>29</sup>


- There have been relatively few adverse events reported after HPV. Commonly reported symptoms include injection-site reactions such as pain, redness and swelling, as well as dizziness, fainting, nausea, and headache.<sup>30</sup> A review of eight years of post-licensure follow-up studies of Gardasil® worldwide found no serious health risks associated with the vaccine.<sup>31</sup>

- Prevalence of HPV in the U.S. has decreased significantly since the vaccine was made available, falling from 11.5% to 4.3% among girls ages 14 to 19 between 2003-2006 and 2009-2012. In 2009-2012, half of this group had received at least one-dose of the vaccine.<sup>32</sup>

## VACCINE FINANCING

There are multiple sources of private and public financing that assure that nearly all children and young adults in the U.S. have coverage for the HPV vaccine. Many of the financing entities base their coverage on the recommendations of the CDC's Advisory Community for Immunization Practices (ACIP) an independent body of experts that reviews the immunization recommendations for the U.S. population.

- The Affordable Care Act (ACA) requires most [private insurance plans](#) to cover some recommended preventive services and ACIP recommended immunizations without consumer cost-sharing. Plans must cover the HPV vaccine for the recommended age groups of males and females, Pap tests, and HPV testing for women are all covered under this policy.
- Individuals who obtain insurance through the ACA marketplace are also covered for the HPV vaccine, Pap Tests, and HPV DNA testing without cost-sharing.



## PUBLIC FINANCING

- **Medicaid** – The VFC pays for vaccinations for all children through age 18 with Medicaid. Women and men ages 19 and 20 with Medicaid are eligible for Medicaid coverage of all ACIP-recommended vaccines as an Early and Periodic Screening Diagnosis and Treatment program (EPSDT) service.<sup>33</sup> Adults who qualified for Medicaid as a result of the ACA expansion are covered for the vaccine, HPV testing, as well as pap tests without cost-sharing in states that expanded coverage. For adults 21 and older who qualify for Medicaid through other eligibility pathways, vaccine coverage is an optional benefit and is decided on a state-by-state basis. As of 2015, at least [39 states and DC](#) reported covering the vaccine for women.
- **Vaccines for Children (VFC) Program** – This federally-financed program pays for vaccines recommended by the ACIP for children ages 18 and under who are either Medicaid-eligible, uninsured, American Indian or Alaska Native, or underinsured. ACIP recommended VFC coverage for Gardasil 9 in February 2015 for males and females ages 9 through 18.
- **Immunization Grant Program (Section 317 of the Public Health Service Act)** – Provides grants to states and local agencies to help extend the availability of vaccines to uninsured adults in the United States. Merck has also established assistance programs to provide free vaccines to uninsured low-income adults in the United States.<sup>34</sup> These funds are often directed towards meeting the needs of priority populations, such as underinsured children and uninsured adults.<sup>35</sup>
- **Children’s Health Insurance Program (CHIP)** – State CHIP programs that are separate from their Medicaid programs must cover ACIP-recommended vaccines for beneficiaries since they are not eligible for coverage under the federal VFC.<sup>36</sup>

\*\*\*

The HPV vaccine has been available in the U.S. for several years and uptake is rising. Since its introduction 11 years ago, the vaccine covers more strains of HPV, its use has been extended to males as well as females, the dosage has dropped from three to two shots, and the cost is fully covered by private insurance and public programs. With these improvements, the vaccine holds the promise to safely and dramatically reduce rate of and prevent many kinds of cancers that have long been responsible for the deaths of women and men across the United States and the world.

# Endnotes

- <sup>1</sup> Centers for Disease Control and Prevention (CDC). [Genital HPV Infection—CDC Factsheet](#). July 2017.
- <sup>2</sup> CDC. [Genital HPV Infection—CDC Factsheet](#). July 2017.
- <sup>3</sup> McQuillan G, et al. [Prevalence of HPV in Adults Aged 18-69: United States 2011-2014](#). NCHS Data Brief, No. 280. April 2017.
- <sup>4</sup> CDC. (2014). [Human Papillomavirus Vaccination: Recommendations of the Advisory Committee on Immunization Practices \(ACIP\)](#). *MMWR* 63(rr05).
- <sup>5</sup> CDC. [Genital HPV Infection—CDC Factsheet](#). July 2017.
- <sup>6</sup> National Cancer Institute. [Cancer Stat Facts: Cervix Uteri Cancer](#). Accessed June 2017.
- <sup>7</sup> U.S. Preventive Services Task Force. [Cervical Cancer: Screening](#). March 2012.
- <sup>8</sup> Downs LS, et al. (2008). [The disparity of cervical cancer in diverse populations](#). *Gynecologic Oncology* 109(2).
- <sup>9</sup> National Center for Health Statistics. [Health, United States, 2015: With Special Feature on Racial and Ethnic Health Disparities](#). May 2016.
- <sup>10</sup> Office of Minority Health. (2009). Eliminate disparities in cancer screening & management.
- <sup>11</sup> CDC. [How Many Cancers are Linked with HPV Each Year?](#). March 2017.
- <sup>12</sup> CDC. (2016). [Human Papillomavirus-Associated Cancers- United States, 2008-2012](#). *MMWR* 65(26).
- <sup>13</sup> Saraiya M, et al. (2015). US Assessment of HPV Types in Cancers: Implications for Current and 9-Valent HPV Vaccines. *JNCI Journal of the National Cancer Institute*, 107(6).
- <sup>14</sup> National Cancer Institute. [HPV and Cancer](#). November 2, 2016., National Cancer Institute. [Cancer Stat Facts: Anal Cancer](#). Accessed June 2017.
- <sup>15</sup> CDC. [HPV and Men- Fact Sheet](#). July 2017.
- <sup>16</sup> CDC. [HPV Vaccine Information for Clinicians](#). December 2016.
- <sup>17</sup> Joura E, et al. (2007). [Efficacy of a Quadrivalent Prophylactic Human Papillomavirus \(types 6, 11, 16, and 18\) L1 virus-like-particle vaccine against high-grade vulval and vaginal lesions: A combined analysis of three randomized clinical trials](#). *The Lancet*, 369(9574).
- <sup>18</sup> Garland, S., et al. (2007). [Quadrivalent vaccine against Human Papillomavirus to prevent anogenital diseases](#). *NEJM*, 356(19).
- <sup>19</sup> Meites E, Kempe A, Markowitz LE. (2016). [Use of a 2-Dose Schedule for Human Papillomavirus Vaccination—Updated Recommendations of the Advisory Committee on Immunization Practices](#). CDC. *MMRW* 65(49).
- <sup>20</sup> Ibid.
- <sup>21</sup> CDC. [HPV Vaccine for Preteens and Teens](#). June 2017.
- <sup>22</sup> Westra, TA, et al. (2011). [Until which age should women be vaccinated against HPV infection? Recommendation based on cost-effectiveness analysis](#). *Journal of Infectious Diseases*, 204(3).
- <sup>23</sup> National Health and Medical Research Council. [The Australian Immunisation Handbook](#). Department of Health and Ageing, Australian Government; 2017.
- <sup>24</sup> CDC. [Human Papillomavirus \(HPV\), Questions and Answers](#). November 2016.
- <sup>25</sup> HPV UTD includes those with  $\geq 3$  doses, and those with 2 doses when the first HPV Vaccine dose was initiated before age 15 years and time between the first and the second dose was at least 5 months minus 4 days.
- <sup>26</sup> CDC. (2017). [National, Regional, State, Selected Local Area Vaccination Coverage Among Adolescents Aged 13-17 Years—United States, 2016](#). *MMRW* 66(33).
- <sup>27</sup> Ibid.
- <sup>28</sup> National Conference of State Legislatures. [HPV Vaccine: State Legislation and Statues](#). July 2017.
- <sup>29</sup> State of Rhode Island Department of Health. [Immunization Information for Schools and Child Care Workers](#). Accessed August 2017.
- <sup>30</sup> CDC. [HPV \(Human Papillomavirus\) VIS](#). December 2016.
- <sup>31</sup> Vichnin M, et al. (2015). [An Overview of Quadrivalent Human Papillomavirus Vaccine Safety: 2006 to 2015](#). *The Pediatric Infectious Disease Journal*, 34(9).
- <sup>32</sup> Markowitz LE, et al. (2016). [Prevalence of HPV After Introduction of the Vaccination Program in the United States](#). *Pediatrics*, 137(2).
- <sup>33</sup> Sebelius K. [Report to Congress on Preventive Services and Obesity-related Services Available to Medicaid Enrollees](#). 2014.
- <sup>34</sup> National Cancer Institute. [Human Papillomavirus \(HPV\) Vaccines](#). November 2016.
- <sup>35</sup> CDC. [Questions Answered on Vaccines Purchased with 317 funds](#). February 2016.
- <sup>36</sup> Association of Maternal and Child Health Programs. (2007). The HPV vaccine: background, coverage, & benefits.